

INEQUALITIES FOR THE POLAR DERIVATIVE OF A POLYNOMIAL

M. Shakeri, M. Bidkham and M.Eshaghi Gordji

Department of Mathematics, Faculty of Natural Sciences
Semnan University, Semnan, Iran.

E-mail: mbidkham@semnan.ac.ir, madjid.eshaghi@gmail.com

Abstract

In this paper we obtain new results concerning maximum modulus of the polar derivative of a polynomial with restricted zeros. Our results generalize and refine upon the results of Aziz and Shah [An integral mean estimate for polynomial, Indian J. Pure Appl. Math. 28 (1997) 1413–1419] and Gardner, Govil and Weems [Some result concerning rate of growth of polynomials, East J. Apporox. 10(2004) 301–312].

1. INTRODUCTION AND STATEMENT OF RESULTS

The problems in the analytic theory of polynomials concerning derivative of the polynomials have been frequently investigated. Over many decades, a large number of research papers, e.g, [1, 2, 3, 4, 5, 9, 11] have been published.

If $p(z) = \sum_{m=0}^n a_m z^m$ is a polynomial of degree n , then

$$\max_{|z|=1} |p'(z)| \leq n \max_{|z|=1} |p(z)|. \quad (1)$$

The above inequality, which is an immediate consequence of Bernstein's inequality on the derivative of a trigonometric polynomial is best possible with equality holding for the polynomial $p(z) = \lambda z^n$, λ being a complex number.

2000 Mathematics Subject Classification. 30A10, 30C15, 26C10, 12D10, 30D15.

Key words and phrases: Polynomials, zeros of polynomials, Polar Derivative.

It is noted that in (1.1) equality hold if and only if $p(z)$ has all its zeros at the origin and so it is natural to seek improvements under appropriate assumptions on the zeros of $p(z)$.

If $p(z)$ having no zeros in $|z| < 1$, then the above inequality can be replaced by

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{2} \max_{|z|=1} |p(z)|. \quad (2)$$

Inequality (2) was conjectured by Erdos and later proved by Lax [10]. On the other hand, it was shown by Turan [12] that if all the zeros of $p(z)$ lie in $|z| < 1$, then

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{2} \max_{|z|=1} |p(z)|. \quad (3)$$

The above inequality was generalized by Govil [9]. Who proved that if $p(z)$ is a polynomial of degree n having all its zeros in $|z| < k$, then for $k \leq 1$

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1+k} \max_{|z|=1} |p(z)|, \quad (4)$$

and for $k \geq 1$

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|. \quad (5)$$

Both the above inequalities are best possible with equality in (4) holding for $p(z) = (z+k)^n$, while in (5) the equality holds for the polynomial $p(z) = z^n + k^n$. As an extension of (2) was shown by Malik [11] that, if $p(z) \neq 0$ in $|z| < k, k \geq 1$, then

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k} \max_{|z|=1} |p(z)|. \quad (6)$$

Equality in (6) holds for $p(z) = (z+k)^n$.

By considering a more general class of polynomials $p(z) = a_0 + \sum_{\nu=\mu}^n a_\nu z^\nu$, $1 \leq \mu \leq n$, not vanishing in $|z| < k, k > 0$, then for $0 < r \leq R \leq k$, inequality (6) is generalized by Aziz and Shah [4] by proving

$$\max_{|z|=R} |p'(z)| \leq nR^{\mu-1} \frac{(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} [\max_{|z|=r} |p(z)| - \min_{|z|=k} |p(z)|]. \quad (7)$$

Equality in (7) holds for $p(z) = (z^\mu + k^\mu)^{\frac{n}{\mu}}$ where n is a multiple of μ . On the other hand, for the class of polynomial $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, of degree n having all its zeros in $|z| \leq k, k \leq 1$, Aziz and Shah [2] proved

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1+k^\mu} \left\{ \max_{|z|=1} |p(z)| + \frac{1}{k^{n-\mu}} \min_{|z|=k} |p(z)| \right\}. \quad (8)$$

Let $D_\alpha\{p(z)\}$ denote the polar derivative of the polynomial $p(z)$ of degree n with respect to the point α , then

$$D_\alpha\{p(z)\} = np(z) + (\alpha - z)p'(z).$$

The polynomial $D_\alpha\{p(z)\}$ is of degree at most $n - 1$ and it generalized the ordinary derivative in the sense that

$$\lim_{\alpha \rightarrow \infty} \left[\frac{D_\alpha\{p(z)\}}{\alpha} \right] = p'(z)$$

Dewan [7] extended (8) to the polar derivative of a polynomial and proved the following.

Theorem A Let $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, be a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$. Then for every real or complex number α with $|\alpha| \geq k^\mu$, we have

$$\begin{aligned} \max_{|z|=1} |D_\alpha p(z)| &\geq \frac{n(|\alpha| - k^\mu)}{1 + k^\mu} \max_{|z|=1} |p(z)| + n \left(\frac{|\alpha| + 1}{k^{n-\mu}(1 + k^\mu)} \right) m \\ &\quad n \left(\frac{k^\mu - A_\mu}{1 + k^\mu} \right) \max_{|z|=1} |p(z)| + \frac{n(A_\mu - k^\mu)}{k^n(1 + k^\mu)} m, \end{aligned} \quad (9)$$

where $m = \min_{|z|=k} |p(z)|$ and

$$A_\mu = \frac{n(|a_n| - \frac{m}{k^n})k^{2\mu} + \mu|a_{n-\mu}|k^{\mu-1}}{n(|a_n| - \frac{m}{k^n})k^{\mu-1} + \mu|a_{n-\mu}|}.$$

Dividing both sides of inequality (9) by $|\alpha|$ and let $|\alpha| \rightarrow \infty$ we get (8).

As an extension of (6) to the polar derivative of a polynomial, we have the following result due to Dewan [6].

Theorem B If $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq \frac{n}{1 + k^\mu} \{ (|\alpha| + k^\mu) \max_{|z|=1} |p(z)| - (|\alpha| - 1)m \} \quad (10)$$

where $m = \min_{|z|=k} |p(z)|$.

The above theorem is an extension of a result of Aziz [1] and for $\mu = 1$, it reduces to a result of Aziz and Shah [2].

In this paper, we first obtain the following generalization of inequality (8) which is also a refinement of inequality (9).

Theorem 1. Let $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, be a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$ and α is any real or complex number with $|\alpha| \geq \frac{k^\mu}{R^{\mu-1}}$, then for $rR \geq k^2$ and $r \leq R$, we have

$$\begin{aligned}
\max_{|z|=R} |D_\alpha p(z)| &\geq n(R^{\mu-1}|\alpha| - k^\mu) \frac{(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\
&+ \frac{n(R^{\mu-1}|\alpha| + R^\mu)}{k^{n-\mu}(R^\mu + k^\mu)} \min_{|z|=k} |p(z)| \\
&+ n(k^\mu - R^\mu A'_\mu) \frac{(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\
&+ \frac{nR^n(R^\mu A'_\mu - k^\mu)}{k^n(R^\mu + k^\mu)} \min_{|z|=k} |p(z)| \\
&+ \frac{nR^{\mu-1}}{(R^\mu + k^\mu)} (|\alpha| - RA'_\mu) \left[\left(\frac{R}{r}\right)^n - \left(\frac{R^\mu + k^\mu}{r^\mu + k^\mu}\right)^{\frac{n}{\mu}} \right] \min_{|z|=k} |p(z)|, \quad (11)
\end{aligned}$$

where

$$A'_\mu = \frac{n(|a_n| - \frac{m}{k^n}) \frac{k^{2\mu}}{R^\mu} + \mu |a_{n-\mu}| \frac{k^{\mu-1}}{R^{\mu-1}}}{nR(|a_n| - \frac{m}{k^n}) k^{\mu-1} + \mu |a_{n-\mu}|}$$

and $m = \min_{|z|=k} |p(z)|$.

Remark 1. For $R = r = 1$ theorem 1 reduces to (9).

Remark 2. Dividing the two sides of (11) by $|\alpha|$, letting $|\alpha| \rightarrow \infty$, we get

$$\begin{aligned}
\max_{|z|=R} |p'(z)| &\geq \frac{nR^{\mu-1}(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| + \frac{nR^{\mu-1}}{k^{n-\mu}(R^\mu + k^\mu)} \min_{|z|=k} |p(z)| \\
&+ \frac{nR^{\mu-1}}{(R^\mu + k^\mu)} \left[\left(\frac{R}{r}\right)^n - \left(\frac{R^\mu + k^\mu}{r^\mu + k^\mu}\right)^{\frac{n}{\mu}} \right] \min_{|z|=k} |p(z)|. \quad (12)
\end{aligned}$$

This includes inequality (8) as special case. The following result immediately follows by taking $k = 1$ in theorem 1.

Corollary 1. If $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, be a polynomial of degree n having all its zeros in $|z| \leq 1$ and α is any real or complex number α with

$|\alpha| \geq \frac{1}{R^{\mu-1}}$, then for $rR \geq 1$ and $r \leq R$, we have

$$\begin{aligned}
\max_{|z|=R} |D_\alpha p(z)| &\geq \frac{n(R^{\mu-1}|\alpha| + R^\mu)}{R^\mu + 1} \min_{|z|=1} |p(z)| \\
&\quad + n(R^{\mu-1}|\alpha| - 1) \frac{(R^\mu + 1)^{\frac{n}{\mu}-1}}{(r^\mu + 1)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\
&\quad + n(1 - R^\mu A''_\mu) \frac{(R^\mu + 1)^{\frac{n}{\mu}-1}}{(r^\mu + 1)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\
&\quad + \frac{nR^n(R^\mu A''_\mu - 1)}{(R^\mu + 1)} \min_{|z|=1} |p(z)| \\
&\quad + \frac{nR^{\mu-1}}{(R^\mu + 1)} (|\alpha| - RA''_\mu) \left[\left(\frac{R}{r}\right)^n - \frac{(R^\mu + 1)^{\frac{n}{\mu}}}{(r^\mu + 1)} \right] \min_{|z|=1} |p(z)|, \quad (13)
\end{aligned}$$

where

$$A''_\mu = \frac{\frac{n}{R^\mu}(|a_n| - m) + \frac{\mu}{R^{\mu-1}}|a_{n-\mu}|}{nR(|a_n| - m) + \mu|a_{n-\mu}|}$$

and $m = \min_{|z|=1} |p(z)|$.

We next prove the following result which is a generalization of the inequality (7), and result due to Gardner, Govil and Weems [8].

Theorem 2. If $p(z) = a_0 + \sum_{\nu=m}^n a_\nu z^\nu$, $1 \leq m \leq n$, be a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$ for $0 \leq r \leq R \leq k$, then for every real or complex number α with $|\alpha| \geq R$,

$$\begin{aligned}
\max_{|z|=R} |D_\alpha p(z)| &\leq \frac{n(R^{m-1}|\alpha| + k^m)}{R^m + k^m} \left[\left(\frac{R^m + k^m}{r^m + k^m}\right)^{\frac{n}{m}} \max_{|z|=r} |p(z)| \right. \\
&\quad \left. - \left(\left(\frac{R^m + k^m}{r^m + k^m}\right)^{\frac{n}{m}} - 1\right) \min_{|z|=k} |p(z)| \right] \\
&\quad - \frac{n(R^{m-1}|\alpha| - R^m)}{R^m + k^m} \min_{|z|=k} |p(z)|. \quad (14)
\end{aligned}$$

Equality in (14) holds for $p(z) = (z^m + k^m)^{\frac{n}{m}}$.

Remark 3. For $R = r = 1$ theorem 2 reduces to theorem B.

Remark 4. Dividing the two sides of (14) by $|\alpha|$ and letting $|\alpha| \rightarrow \infty$, we have the following inequality, which is an improvement as well as a generalization of a result proved by Bidkham and Dewan [5]

$$\max_{|z|=R} |p'(z)| \leq \frac{nR^{m-1}(R^m + k^m)^{\frac{n}{m}-1}}{(r^m + k^m)^{\frac{n}{m}}} \left\{ \max_{|z|=r} |p(z)| - \min_{|z|=k} |p(z)| \right\}.$$

The result is best possible and equality holds for the polynomial $p(z) = (z^m + k^m)^{\frac{n}{m}}$ where n is a multiple of m .

2. LEMMAS

For the proofs of these theorems we need the following lemmas.

Lemma 2.1. If $p(z) = a_0 + \sum_{\nu=m}^n a_\nu z^\nu$, $1 \leq m \leq n$, is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k$, $k > 0$, then for $0 \leq r \leq R \leq k$,

$$\max_{|z|=r} |p(z)| \geq \left(\frac{r^m + k^m}{R^m + k^m}\right)^{\frac{n}{m}} \max_{|z|=R} |p(z)| + \left[1 - \left(\frac{r^m + k^m}{R^m + k^m}\right)^{\frac{n}{m}}\right] \min_{|z|=k} |p(z)|. \quad (15)$$

Here the result is best possible and equality holds for the polynomial $p(z) = (z^m + k^m)^{\frac{n}{m}}$ where n is multiple of m . This result is due to Dewan [7].

Lemma 2.2 . If $p(z) = a_0 + \sum_{\nu=\mu}^n a_\nu z^\nu$, $1 \leq \mu \leq n$, is a polynomial of degree n having no zeros in $|z| \leq k$, $k \geq 1$, then for every real or complex number α with $|\alpha| \geq 1$

$$\max_{|z|=1} |D_\alpha p(z)| \leq \frac{n}{1 + k^\mu} \{(|\alpha| + k^\mu) \max_{|z|=1} |p(z)| - (|\alpha| - 1)m\},$$

where $m = \min_{|z|=k} |p(z)|$.

This result is due to [6].

3. PROOF OF THE THEOREMS

Proof of theorem 1. By hypothesis the polynomial $p(z) = a_n z^n + \sum_{\nu=\mu}^n a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, has all its zeros in $|z| \leq k$, where $k \leq 1$, therefore it follows that $F(z) = P(Rz)$ has all its zeros in $|z| \leq \frac{k}{R}$, where $\frac{k}{R} \leq 1$ and $|\alpha| \geq (\frac{k}{R})^\mu$. Applying inequality (9) to the polynomial $F(z)$ we get

$$\begin{aligned} \max_{|z|=1} |D_\alpha F(z)| &\geq \frac{n(|\alpha| - \frac{k^\mu}{R^\mu})}{(1 + \frac{k^\mu}{R^\mu})} \max_{|z|=1} |F(z)| \\ &\quad + n \left(\frac{|\alpha| + 1}{(\frac{k}{R})^{n-\mu} (1 + \frac{k^\mu}{R^\mu})} \right) \min_{|z|=k/R} |F(z)| \\ &\quad + \frac{n((\frac{k}{R})^\mu - A'_\mu)}{(1 + (\frac{k}{R})^\mu)} \max_{|z|=1} |F(z)| + \frac{n(A'_\mu - \frac{k^\mu}{R^\mu})}{(\frac{k}{R})^n (1 + (\frac{k}{R})^\mu)} \min_{|z|=k/R} |F(z)|, \end{aligned} \quad (16)$$

where

$$A'_\mu = \frac{n(|a_n| - \frac{m}{k^n}) \frac{k^{2\mu}}{R^\mu} + \mu |a_{n-\mu}| \frac{k^{\mu-1}}{R^{\mu-1}}}{nR(|a_n| - \frac{m}{k^n}) k^{\mu-1} + \mu |a_{n-\mu}|}$$

and $m = \min_{|z|=k} |p(z)|$,
which is equivalent to

$$\begin{aligned} \max_{|z|=1} |np(Rz) + (\alpha - z)Rp'(Rz)| &\geq \frac{n(|\alpha|R^\mu - k^\mu)}{R^\mu + k^\mu} \max_{|z|=1} |p(Rz)| \\ &+ \frac{nR^n(|\alpha| + 1)}{K^{n-\mu}(R^\mu + K^\mu)} \min_{|z|=k/R} |p(Rz)| + n \frac{(K^\mu - R^\mu A'_\mu)}{(R^\mu + k^\mu)} \max_{|z|=1} |p(Rz)| \\ &+ \frac{nR^n(R^\mu A'_\mu - k^\mu)}{k^n(R^\mu + k^\mu)} \min_{|z|=k/R} |p(Rz)|, \end{aligned}$$

which gives

$$\begin{aligned} \max_{|z|=R} |np(z) + (\alpha R - z)p'(z)| &\geq \frac{n(|\alpha R|R^{\mu-1} - k^\mu)}{R^\mu + k^\mu} \max_{|z|=R} |p(z)| \\ &+ \frac{n(|\alpha R|R^{n-1} + R^n)}{K^{n-\mu}(R^\mu + K^\mu)} \min_{|z|=k} |p(z)| + n \left(\frac{K^\mu - R^\mu A'_\mu}{R^\mu + k^\mu} \right) \max_{|z|=R} |p(z)| \\ &+ \frac{nR^n(R^\mu A'_\mu - k^\mu)}{k^n(R^\mu + k^\mu)} \min_{|z|=k} |p(z)|. \end{aligned} \quad (17)$$

On other hand, since $p(z)$ has all its zeros in $|z| < k, k > 0$ therefore it follows that $q(z) = z^n \overline{p(\frac{1}{z})} \neq 0$ for $|z| < 1/k$. Applying inequality (15) to $q(z)$, we get

$$\max_{|z|=1/R} |q(z)| \geq \left(\frac{(\frac{1}{k})^\mu + (\frac{1}{R})^\mu}{(\frac{1}{k})^\mu + (\frac{1}{r})^\mu} \right)^{\frac{n}{\mu}} \max_{|z|=1/r} |q(z)| + \left[1 - \left(\frac{(\frac{1}{k})^\mu + (\frac{1}{R})^\mu}{(\frac{1}{k})^\mu + (\frac{1}{r})^\mu} \right)^{\frac{n}{\mu}} \right] \min_{|z|=1/k} |q(z)|$$

, which gives

$$\max_{|z|=R} |p(z)| \geq \left(\frac{R^\mu + k^\mu}{r^\mu + k^\mu} \right)^{\frac{n}{\mu}} \max_{|z|=r} |p(z)| + \left[\left(\frac{R}{r} \right)^n - \left(\frac{R^\mu + k^\mu}{r^\mu + k^\mu} \right)^{\frac{n}{\mu}} \right] \min_{|z|=k} |p(z)|. \quad (18)$$

From (17) and (18), we have

$$\begin{aligned} \max_{|z|=R} |D_{R\alpha}p(z)| &\geq n(|\alpha R|R^{\mu-1} - k^\mu) \frac{(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\ &+ \frac{n(R^{n-1}|\alpha R| + R^n)}{k^{n-\mu}(R^\mu + k^\mu)} \min_{|z|=k} |p(z)| \\ &+ n(k^\mu - R^\mu A'_\mu) \frac{(R^\mu + k^\mu)^{\frac{n}{\mu}-1}}{(r^\mu + k^\mu)^{\frac{n}{\mu}}} \max_{|z|=r} |p(z)| \\ &+ \frac{nR^n(R^\mu A'_\mu - k^\mu)}{k^n(R^\mu + k^\mu)} \min_{|z|=k} |p(z)| \\ &+ \frac{nR^{\mu-1}}{(R^\mu + k^\mu)} (|\alpha R| - RA'_\mu) \left[\left(\frac{R}{r} \right)^n - \left(\frac{R^\mu + k^\mu}{r^\mu + k^\mu} \right)^{\frac{n}{\mu}} \right] \min_{|z|=R} |p(z)|, \end{aligned}$$

which is equivalent to (11).

Proof of theorem 2. By hypothesis the polynomial $p(z) = a_0 + \sum_{\nu=m}^n a_\nu z^\nu$, $1 \leq m \leq n$, having no zeros in $|z| < k$, where $k \geq 1$, therefore it follows that $F(z) = p(Rz)$ having no zeros in $|z| \leq \frac{k}{R}$, where $\frac{k}{R} \geq 1$. Hence using lemma 2 for $|\alpha| \geq 1$. we have

$$\max_{|z|=1} |D_\alpha F(z)| \leq \frac{n}{1 + (\frac{k}{R})^m} \{ (|\alpha| + (\frac{k}{R})^m) \max_{|z|=1} |F(z)| - (|\alpha| - 1) \min_{|z|=k/R} |F(z)| \},$$

which is equivalent to

$$\max_{|z|=1} |np(Rz) + (\alpha - z)Rp'(Rz)| \leq \frac{n}{1 + (\frac{k}{R})^m} \{ (|\alpha| + (\frac{k}{R})^m) \max_{|z|=1} |p(Rz)| - (|\alpha| - 1) \min_{|z|=k/R} |p(Rz)| \}.$$

Replacing Rz by z we get

$$\begin{aligned} \max_{|z|=R} |D_{R\alpha} p(z)| &\leq \frac{n(R^{m-1}|\alpha R| + k^m)}{R^m + k^m} \max_{|z|=R} |p(z)| \\ &\quad - \frac{n(R^{m-1}|R\alpha| - R^m)}{(R^m + k^m)} \min_{|z|=k} |p(z)| \end{aligned}$$

for $R \geq 1$

or

$$\begin{aligned} \max_{|z|=R} |D_\alpha p(z)| &\leq \frac{n(R^{m-1}|\alpha| + k^m)}{R^m + k^m} \max_{|z|=R} |p(z)| \\ &\quad - \frac{n(R^{m-1}|\alpha| - R^m)}{(R^m + k^m)} \min_{|z|=k} |p(z)|. \end{aligned} \tag{19}$$

By combining (19) and (15) we get (11).

References

- [1] A. AZIZ, Inequalities for the polar derivative of a polynomial, J. Approx. Theory 55 (1988) 183–193.
- [2] A. AZIZ AND W.M. SHAH, An integral mean estimate for polynomial, Indian J. Pure Appl. Math. 28 (1997) 1413–1419.
- [3] A. AZIZ AND W.M. SHAH, Inequalities for polar derivative of a polynomial, Indian J. Pure Appl. Math. 29 (1998) 163–173.
- [4] A. AZIZ AND W.M. SHAH, Inequalities for a polynomial and its derivative, Math. Inequal. Appl. 7(3)(2004) 379–391.

- [5] M. BIDKHAM AND K.K. DEWAN, Inequalities for polynomial and its derivative, J. Math. Anal. Appl. 166 (1992) 319–324.
- [6] K.K. DEWAN AND NARESH SINGH and ABDULLAH MIR, Extensions of some polynomial, inequalities to the polar derivative, J. Math. Anal. Appl. 352 (2009), no. 2, 807–815.
- [7] K.K. DEWAN, R.S. YADAV AND M.S.PUKHTA, Inequalities for a polynomial and its derivative, Math. Inequal. Appl. 2(2) (1999) 203–205.
- [8] R.B. GARDNER, N.K. GOVIL AND A.WEEMS, Some result concerning rate of growth of polynomials, East J. Apporox. 10(2004) 301–312.
- [9] N.K. GOVIL, On the derivative of a polynomial, Proc. Amer. Math. Soc. 41(1973) 643–546.
- [10] P.D. LAX, Proof of a conjecture of P.Erdos on the derivative of a polynomial, Bull. Amer. Math. Soc. 50 (1944) 509–513.
- [11] M.A. MALIK, On the derivative of a polynomial, J. Londen Math. Soc. 1(1969) 57–60.
- [12] P. TURAN, Uberdie ableting von polynoment, Compos. Math. 7(1939) 82–95.